

Exhibit 5



Standards of Care

for the Health of Transsexual, Transgender, and Gender Nonconforming People

The World Professional Association for Transgender Health

7th Version¹ | www.wpath.org

¹ This is the seventh version of the Standards of Care. The original SOC were published in 1979. Previous revisions were in 1980, 1981, 1990, 1998, and 2001.

for gender and sexual diversity and that eliminate prejudice, discrimination, and stigma. WPATH is committed to advocacy for these changes in public policies and legal reforms.

The Standards of Care Are Flexible Clinical Guidelines

The SOC are intended to be flexible in order to meet the diverse health care needs of transsexual, transgender, and gender nonconforming people. While flexible, they offer standards for promoting optimal health care and guiding the treatment of people experiencing gender dysphoria – broadly defined as discomfort or distress that is caused by a discrepancy between a person's gender identity and that person's sex assigned at birth (and the associated gender role and/or primary and secondary sex characteristics) (Fisk, 1974; Knudson, De Cuypere, & Bockting, 2010b).

As for all previous versions of the SOC, the criteria put forth in this document for hormone therapy and surgical treatments for gender dysphoria are clinical guidelines; individual health professionals and programs may modify them. Clinical departures from the SOC may come about because of a patient's unique anatomic, social, or psychological situation; an experienced health professional's evolving method of handling a common situation; a research protocol; lack of resources in various parts of the world; or the need for specific harm reduction strategies. These departures should be recognized as such, explained to the patient, and documented through informed consent for quality patient care and legal protection. This documentation is also valuable for the accumulation of new data, which can be retrospectively examined to allow for health care – and the SOC – to evolve.

The SOC articulate standards of care but also acknowledge the role of making informed choices and the value of harm reduction approaches. In addition, this version of the SOC recognizes and validates various expressions of gender that may not necessitate psychological, hormonal, or surgical treatments. Some patients who present for care will have made significant self-directed progress towards gender role changes, transition, or other resolutions regarding their gender identity or gender dysphoria. Other patients will require more intensive services. Health professionals can use the SOC to help patients consider the full range of health services open to them, in accordance with their clinical needs and goals for gender expression.

Among adolescents who are referred to gender identity clinics, the number considered eligible for early medical treatment – starting with GnRH analogues to suppress puberty in the first Tanner stages – differs among countries and centers. Not all clinics offer puberty suppression. If such treatment is offered, the pubertal stage at which adolescents are allowed to start varies from Tanner stage 2 to stage 4 (Delemarre-van de Waal & Cohen-Kettenis, 2006; Zucker et al., in press). The percentages of treated adolescents are likely influenced by the organization of health care, insurance aspects, cultural differences, opinions of health professionals, and diagnostic procedures offered in different settings.

Inexperienced clinicians may mistake indications of gender dysphoria for delusions. Phenomenologically, there is a qualitative difference between the presentation of gender dysphoria and the presentation of delusions or other psychotic symptoms. The vast majority of children and adolescents with gender dysphoria are not suffering from underlying severe psychiatric illness such as psychotic disorders (Steensma, Biemond, de Boer, & Cohen-Kettenis, published online ahead of print January 7, 2011).

It is more common for adolescents with gender dysphoria to have co-existing internalizing disorders such as anxiety and depression, and/or externalizing disorders such as oppositional defiant disorder (de Vries et al., 2010). As in children, there seems to be a higher prevalence of autistic spectrum disorders in clinically referred, gender dysphoric adolescents than in the general adolescent population (de Vries et al., 2010).

Competency of Mental Health Professionals Working with Children or Adolescents with Gender Dysphoria

The following are recommended minimum credentials for mental health professionals who assess, refer, and offer therapy to children and adolescents presenting with gender dysphoria:

1. Meet the competency requirements for mental health professionals working with adults, as outlined in section VII;
2. Trained in childhood and adolescent developmental psychopathology;
3. Competent in diagnosing and treating the ordinary problems of children and adolescents.

Competency of Mental Health Professionals Working with Adults Who Present with Gender Dysphoria

The training of mental health professionals competent to work with gender dysphoric adults rests upon basic general clinical competence in the assessment, diagnosis, and treatment of mental health concerns. Clinical training may occur within any discipline that prepares mental health professionals for clinical practice, such as psychology, psychiatry, social work, mental health counseling, marriage and family therapy, nursing, or family medicine with specific training in behavioral health and counseling. The following are recommended minimum credentials for mental health professionals who work with adults presenting with gender dysphoria:

1. A master's degree or its equivalent in a clinical behavioral science field. This degree or a more advanced one should be granted by an institution accredited by the appropriate national or regional accrediting board. The mental health professional should have documented credentials from a relevant licensing board or equivalent for that country.
2. Competence in using the *Diagnostic Statistical Manual of Mental Disorders* and/or the *International Classification of Diseases* for diagnostic purposes.
3. Ability to recognize and diagnose co-existing mental health concerns and to distinguish these from gender dysphoria.
4. Documented supervised training and competence in psychotherapy or counseling.
5. Knowledgeable about gender nonconforming identities and expressions, and the assessment and treatment of gender dysphoria.
6. Continuing education in the assessment and treatment of gender dysphoria. This may include attending relevant professional meetings, workshops, or seminars; obtaining supervision from a mental health professional with relevant experience; or participating in research related to gender nonconformity and gender dysphoria.

In addition to the minimum credentials above, it is recommended that mental health professionals develop and maintain cultural competence to facilitate their work with transsexual, transgender, and gender nonconforming clients. This may involve, for example, becoming knowledgeable about current community, advocacy, and public policy issues relevant to these clients and their families. Additionally, knowledge about sexuality, sexual health concerns, and the assessment and treatment of sexual disorders is preferred.